# ROTATOR CUFF REPAIR Arthroscopic and Open

# Antoine I. Jabbour, MD www.drjabbour.com

Tulsa Bone and Joint 918-392-1400

Union Pines Surgery Center 918-742-2502

#### Follow-Up Care

Call the doctor's office today to confirm or make an appointment in about 10 days. Your sutures will be removed at your first post-operative visit.

#### **Activity**

Rest today.

Ice the shoulder for 3-5 days. One hour on, one hour off. Keep a very thin towel between skin and ice bag.

CPM or continuous passive motion machine (if provided) use is to begin day after surgery. Use as tolerated although the representative providing the machine will recommend 2 hours three times per day.

Your arm may be numb first 8-36 hours.

Discontinue sling the day after surgery while in seated position only.

Sling must be on when up walking

Do not reach above shoulder level

Pendulum excercises are performed by hanging your arm to the ground while you are bent over the waist. Dangle fingertips to the ground making gentle small clockwise and counterclockwise circles with your hand. Pendulum excercises should be done for 20-60 seconds, 4 times per day.

### **Care of Incision and Dressing**

It is normal to have some bleeding for 48-72 hours. Reinforce dressing if necessary. Remove dressing and shower in 48-72 hours. Change Band-Aids daily.

Do not submerge wound in bath water, pool, lake water, or hot tub.

Keep incisions clean and dry.

Do not put creams or lotions on your wounds.

#### Medications

Resume routine medications.

Take one Aspirin 325 mg daily for next 5 days.

Most patients will receive <u>Oxvcodone</u> medication for pain,

Morphine pills are for breakthrough pain only (first 48 hours only) and Phenergan for nausea.

#### **Driving**

If left shoulder, you may drive on the third post operative day.

If right shoulder, do not drive until after follow up visit. Do not drive if you are taking narcotic pain medication.

## Call Dr Jabbour's Office If:

If medication causes severe uncontrollable nausea.

If you have shortness of breath or severe calf pain, go the emergency room if you cannot contact the office

If you have a temperature greater than 101.5 degrees Farenheit